



## PERSPECTIVES

# Global transmission of multiple-drug resistant *Neisseria gonorrhoeae* strains refractive to cephalosporin treatment

Shu-Ying Li\*

Research and Diagnostic Center, Centers for Disease Control, Taipei, Taiwan

Received 24 February 2012; received in revised form 1 March 2012; accepted 6 March 2012

Recently, in many countries gonorrhea cases are on the rise. The World Health Organization estimated global annual gonorrhea cases of 62 million in 1999. Gonorrhea infection may increase susceptibility to and transmission of human immunodeficiency virus (HIV). In addition, due to sharing of same transmission route and overlapping risk groups, gonorrhea may serve as a risk marker for infection of HIV and other sexually transmitted infections. In Taiwan, according to Statistics of Communicable Diseases and Surveillance of Taiwan-CDC, 2265 new cases were reported for 2010, with an incidence of 9.79 per 100,000 people.

In only a few decades, *Neisseria gonorrhoeae* has rapidly developed resistance to multiple antibiotics, such as sulfanilamide, penicillins, tetracyclines, and fluoroquinolones. Recently, the third generation cephalosporins, or extended-spectrum cephalosporin (ESC), currently the last remaining treatment options in the drug pipeline, were proposed as alternative antibiotics for treatment of gonorrhea. However, susceptibility to all ESCs (oral and injectable) is decreasing rapidly. Alarming, treatment failures with oral ESCs have been reported in many countries. In Taiwan, according to the 2133 isolates (representing 26.7% of the 8,001 reported cases) of gonococci-National Isolate Collection for Epidemiology (G-NICE) surveillance spanning 2008–2011, resistant rates [criteria: European Committee on Antimicrobial Susceptibility Testing (EUCAST) 2011, R:

Minimum Inhibitory Concentration (MIC)  $\geq 0.25$  mg/L and Clinical And Laboratory Standards Institute (CLSI) 2011, S: MIC  $\leq 0.25$  mg/L] to the oral cephalosporin, cefixime was 2.63% for MIC  $\geq 0.25$  mg/L, while for injectable cephalosporin, ceftriaxone was 0% for MIC  $\geq 0.25$  mg/L. The maximum MIC was 0.5 mg/L and 0.19 mg/L for cefixime and ceftriaxone, respectively (our unpublished G-NICE surveillance data).

Molecular epidemiology study employing molecular typing methods such as *N. gonorrhoeae* multiantigen sequence typing (NG-MAST) or serotyping in conjunction with sequencing the *penA* allele encoding penicillin-binding protein 2 and other resistance determinants (*mtrR*, *penB*, *ponA*, and *pilQ*) is helpful in tracing the resistant clones.<sup>1</sup> Recent studies have indicated that most of the gonococcal isolates with decreased susceptibility to the ESC belongs to a few specific major clones. In Taiwan, the majority of cefixime-reduced susceptible strains belong to related clones containing NG-MAST type10 *tpbB* and mosaic types X *penA* allele such as sequence type (ST) 835, 2180, and ST2253. These clones are multidrug resistant (MDR) with a high percentage of resistance rate to penicillin, tetracycline, and ciprofloxacin, and mostly circulate in high-risk homosexual or bisexual sexual networks.<sup>2</sup>

More recently, global rapid spread of ESC-reduced susceptible ST1407 and related STs that share the *tpbB* allele (allele 110), but with different *por* alleles (*por* 908 related) and harbor mosaic *penA* allele types XXXIV have been reported in Japan<sup>3</sup> and in Europe<sup>4–6</sup>, such as England and Wales, Sweden, the Netherlands, Norway,

\* Research and Diagnostic Center, Centers for Disease Control, Number 161, Kun-Yang Street, Taipei, Taiwan.  
E-mail address: [syl@cdc.gov.tw](mailto:syl@cdc.gov.tw).

and Italy. ST1407 is a successful clone and has previously been found in the United States, Australia, and many European countries. The threat is imminent with increasing reports on MDR ST1407 strains render ESC treatment failure in many European countries,<sup>7,8</sup> such as France, Sweden, England and Norway.

The emergence of MDR ST1407 with decreased susceptibility to ESCs, such as ST1407, threatens to erode the last remaining effective agents for treatment of gonorrhea. The rapid spread of these clones might be facilitated by high-risk sexual behavior and should be monitored closely to identify potential treatment failure. Increasing frequency of MDR gonococci with reduced susceptibility to ESC especially in some high-risk groups such as men who have sex with men has been noted.<sup>9</sup> Continuous and global concerted surveillance are paramount to trace the emergence, introduction, and transmission of specific strains as well as the sexual networks they circulate. The finding that these specific clones may transmit across national boundaries and between different sexual networks highlights the need for undertaking intervention as well as risk-reduction consultation in such core-risk groups.<sup>10</sup>

## Acknowledgments

This work was supported by a grant DOH101-DC-2007 from the Centers for Disease Control, Department of Health, Taiwan.

## References

1. Chen PL, Lee HC, Yan JJ, Hsieh YH, Lee NY, Ko NY, et al. High prevalence of mutations in quinolone-resistance-determining regions and *mtrR* loci in polyclonal *Neisseria gonorrhoeae* isolates at a tertiary hospital in Southern Taiwan. *J Formos Med Assoc Taiwan yi zhi* 2010;**109**:120–7.
2. Huang CT, Yen MY, Wong WW, Li LH, Lin KY, Liao MH, et al. Characteristics and dissemination of mosaic penicillin-binding protein 2-harboring multidrug-resistant *Neisseria gonorrhoeae* isolates with reduced cephalosporin susceptibility in northern Taiwan. *Antimicrob Agents Chemother* 2010;**54**:4893–5.
3. Tanaka M, Koga Y, Nakayama H, Kanayama A, Kobayashi I, Saika T, et al. Antibiotic-resistant phenotypes and genotypes of *Neisseria gonorrhoeae* isolates in Japan: identification of strain clusters with multidrug-resistant phenotypes. *Sex Transm Dis* 2011;**38**:871–5.
4. Heymans R, Bruisten SM, Golparian D, Unemo M, de Vries HJ, van Dam AP. Clonally related *Neisseria gonorrhoeae* isolates with decreased susceptibility to the extended-spectrum cephalosporin cefotaxime in Amsterdam, the Netherlands. *Antimicrob Agents Chemother* 2012;**56**:1516–22.
5. Hjeltnes SO, Golparian D, Dedi L, Skutlaberg DH, Haarr E, Christensen A, et al. Phenotypic and genotypic properties of *Neisseria gonorrhoeae* isolates in Norway in 2009: antimicrobial resistance warrants an immediate change in national management guidelines. *Eur J Clin Microbiol Infect Dis: Official Publication of the European Society of Clinical Microbiology* 2012;**31**:1181–6.
6. Carannante A, Prignano G, Cusini M, Matteelli A, Dal Conte I, Ghisetti V, et al. Cefixime and ceftriaxone susceptibility of *Neisseria gonorrhoeae* in Italy from 2006 to 2010. *Clin Microbiol Infect: The Official Publication of the European Society of Clinical Microbiology and Infectious Diseases* 2012;**18**:558–64.
7. Unemo M, Golparian D, Nicholas R, Ohnishi M, Gallay A, Sednaoui P. High-level cefixime- and ceftriaxone-resistant *N. gonorrhoeae* in Europe (France): novel *penA* mosaic allele in a successful international clone causes treatment failure. *Antimicrob Agents Chemother* 2011.
8. Ison CA, Hussey J, Sankar KN, Evans J, Alexander S. Gonorrhoea treatment failures to cefixime and azithromycin in England, 2010. *Euro surveillance: bulletin europeen sur les maladies transmissibles: Eur Commun Dis Bull* 2011;**16**:7–10.
9. Wong WW, Huang CT, Li LH, Chiang CC, Chen BD, Li SY. Molecular epidemiological identification of *Neisseria gonorrhoeae* clonal clusters with distinct susceptibility profiles associated with specific groups at high risk of contracting human immunodeficiency virus and syphilis. *J Clin Microbiol* 2008;**46**:3931–4.
10. Bolan GA, Sparling PF, Wasserheit JN. The emerging threat of untreatable gonococcal infection. *N Engl J Med* 2012;**366**:485–7.